

# Genomic Analysis of BRCA2, and a Comparative Phylogenetic Analysis of BRCA2 in Humans along Different Species: A Candidate for Breast Cancer in Humans

Yaseen N<sup>1</sup>, Nasir MA<sup>2</sup>, Inam H<sup>2</sup>, Mudassar I<sup>2</sup>, Arshad M<sup>2</sup>, Saroosh I<sup>2</sup>, Mirza RF<sup>3</sup>, Naqvi SZ<sup>4</sup> and Mazhar MW<sup>1\*</sup>

<sup>1</sup>Department of Bioinformatics and Biotechnology, Government College University, Faisalabad, Pakistan

<sup>2</sup>Department of medicine and surgery, Hitec-institute of medical sciences taxila cantt, Pakistan

<sup>3</sup>Department of medicine and surgery, Wah Medical College, Wah Cantt, Pakistan

<sup>4</sup>Department of medicine and surgery, Frontiers Medical College, Abbottabad Pakistan

## \*Corresponding author:

Muhammad Waqar Mazhar,  
Department of Bioinformatics and Biotechnology,  
Government College University, Faisalabad,  
Pakistan

Received: 02 Jan 2024

Accepted: 10 Jan 2024

Published: 13 Jan 2024

J Short Name: AJSCCR

## Copyright:

©2024 Mazhar MW, This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially.

## Citation:

Mazhar MW. Genomic Analysis of BRCA2, and a Comparative Phylogenetic Analysis of BRCA2 in Humans along Different Species: A Candidate for Breast Cancer in Humans. *Ame J Surg Clin Case Rep.* 2024; 7(8): 1-7

## Keywords:

BRCA2 gene; Germ-line mutations; Gene structures; Phylogenetic analysis

## 1. Abstract

In DNA homologous recombination repair (HRR) several proteins are encoded by BRCA1 and BRCA2. Any mutation in HRR pathway increase risk of tumor formation as a result of DNA destroying factors like salts of platinum, anthracyclines and novel agents that can damage the DNA homologous repair pathway in the form of inhibitors. Hereditary mutation in BRCA2 gene is linked with germline mutation that contribute to develop breast cancer in women. In current study, phylogenetic analysis of BRCA2 gene in 3 different species is compared to obtain similarities in that gene. Further analysis of gene structure to find exons and introns in these species is done. This proposed study supports the divergence of these 3 species from their common ancestors due to diversity in genes on chromosomes as a result of gene distribution. This current research on BRCA2 gene in humans as compared to other species help us to observe evolutionary changes in that gene through phylogenetic.

## 2. Introduction

BRCA2 gene is considered in human as a tumor suppressor gene (specifically, a responsible caretaker form of a gene), present in all humans and its protein form, is also termed as breast cancer type 2 susceptibility protein, is observed for repairing damages in DNA. BRCA2 are genes that produce such proteins to repair damaged DNA. Every organism get two copies of these genes that is inherited from parents (De Talhouet et al., 2020) [6]. Sometimes BRCA2

genes are also referred as tumor suppressor genes because when they undergo mutation, they develop into cancer. When people inherit such harmful variants, they are at greater risk of several forms of cancer like ovarian and breast cancer (Peixoto et al., 2020) [15]. Harmful variant can be inherited from either parent in BRCA1 and BRCA2 genes. Any mutation in any of these genes carried by child has 50% chances to develop into cancer. Such inherited mutations are termed as germline mutations that pass from parents to offspring. If a woman inherit any mutation in BRCA1 and BRCA2 genes, then the chance of ovarian and breast cancer increase in that woman's lifetime (Bolton et al., 2012) [4].

On chromosome number 13, BRCA2 gene is present. Mutation in this gene is also passed by autosomal pattern of inheritance. As BRCA1 and BRCA2 are responsible to suppressor tumors so they have ability to control cell death and cell division (Lou et al., 2014) [14]. Any mutation in these genes, ultimately enhance the chances of breast cancer in both men and women. Everyone inherit two copies of BRCA2 genes, one from mother and one from father. It has been observed that if a mutation occur in one copy of gene in a person that is inherited from one parent, then still a normal copy of that gene is also present in that individual from another parent (Kemp et al., 2019) [12].

Mutation in both of these genes BRCA1 and BRCA2 cause ovarian, breast and other type of cancers. If a family undergo mutation in BRCA2 gene, then all family members show same type of mu-

tation (Hatano, Tamada, Matsuo, & Hara, 2020) [9]. DNA binding part of BRCA2 gene contain 4 domains and 27 exons that consist of 82.4 kb part of genomic DNA. 11386 bp of mRNA transcript is encoded by BRCA2 gene. Site of transcription is present on 227 bp upstream region of the BRCA2 ORF on first ATG (Jonsson et al., 2019) [11]. Recent research shows that only 5 to 10% cases of breast cancer develop through inherited mutation in BRCA2, while risk of breast cancer or other cancer in BRCA2 mutation depend on several other factors. Men and women carrying a mutation in BRCA2 gene are greater risk of breast, ovarian, prostate, fallopian and pancreatic cancer (Zhao, Wiese, Kwon, Hromas, & Sung, 2019) [22].

### 3. Materials and Methods

#### 3.1. Sequence Retrieval of BRCA2 Gene

Sequence of BRCA2 gene for 3 different species such as Homo sapiens, Pan troglodytes and Gorilla retrieved from NCBI (<https://www.ncbi.nlm.nih.gov/>). Protein sequence of BRCA1 was used as a query from databases. Then BLASTP was used to check the predicted protein in mammalian species including Homo sapiens, Pan troglodytes and Gorilla. Then all the conserved domains were observed for these candidate sequences by using SMART tool (<http://smart.embl-heidelberg.de/>) and pfam domain (PF00170) (<http://pfam.xfam.org/>). Then sequences were checked to find out the BRCA2 gene in all these mammalian species. Then to predict the conserved domains of candidate gene, multiple sequence alignment was performed (Forbes, Fayter, de Kock, & Quek, 2019) [8]. The full-length BRCA2 proteins from humans, chimpanzee and pan troglodytes were aligned in MEGA 7.0.21 and a phylogenetic tree was constructed that showed the sequence alignments.

#### 3.2. Gene Structure and Annotations

The gene structure display server (GSDS) program (<http://gsds.cbi.pku.edu.cn/>) was used to find the intron/exon regions in the gene structure of Homo sapiens, Pan troglodytes and Gorilla. Exact locations on chromosomes for the BRCA2 genes that code for BRCA2 proteins were observed from the GENE database of NCBI (Eccles et al., 2015) [7]. At the short arm telomere position to the long-arm ends of telomere, the genes were organized separately on chromosome number 13 that is linked in ascending order of physical positioning. Data observed on NCBI ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov/)) was used to find all the initial GOS analysis of orthologs in three different mammalian species. The gene annotations of BRCA2 gene in humans, chimpanzee and pan troglodytes were retrieved from BLAST (Burke et al., 2018) [5]. Each sequence of candidate species was used as a query, to separate the BLAST searches and to

form outcomes more authentic and accurate. These sequences help us to find the more close sequences and more distant sequences. After aligning all sequences, repetitive sequences were excluded from the data {Azeem, 2020 #315}.

#### 3.3. Phylogenetic Analysis of BRCA2 Gene:

Phylogenetic analysis of BRCA2 genes done to find the orthologs and homologs between human BRCA2 and two other mammalian BRCA2 genes. Using the BLAST sequence of BRCA2 genes of mammals in MEGA7, a green tree is constructed. Phylogenetic tree shows evolutionary relationship between human and other mammalian species (Théry et al., 2011) [19]. Characterization of BRCA2 gene in three different mammalian species has done by using Clustal Omega 7. By using Maximum likelihood method we inferred the evolutionary history of BRCA2 gene in three different species. This method is based on JTT (Jones Taylor Thornton) model that is alternative model for protein alignment. The evolutionary tree is constructed according to scale with branch lengths showing evolutionary history (Walker et al., 2010) [20].


### 4. Results

#### 4.1. Identification of BRCA2 Gene in Homo Sapiens, Pan Troglodytes and Gorilla

The protein sequence of BRCA2 in candidate species were retrieved from NCBI-BLAST search. In BLAST searches, every sequence was considered as a query to obtain outcomes more precise and accurate. Through these retrieved sequences more possible matches are identified. After aligning all these sequences, repetitive sequences were removed. The signature domain of BRCA2 in these BLAST sequences was observed by using an online tool (SMART <http://smart.embl-heidelberg.de>) and MEME motif that showed the existence of the conserved BRCA2 domains in all candidate species.

#### 4.2. Phylogenetic Analysis of BRCA2 Gene


The characteristics of the evolutionary relationship between BRCA2 from Homo sapiens, Pan troglodytes and Gorilla gorilla was confirmed by the alignment of gene BRCA2 from all these species by using MEGA 7.0.21 to create rooted tree via the maximum likelihood (ML) method. Gene was found in all three candidate species. The following analyses of conserved motifs and structure of genes reinforced the characterization of BRCA2 in all species (Pfeffer, Ho, & Singh, 2017) [16]. The SMART tool analysis manifested that most of the proteins of BRCA2 genes is conserved in all three candidate species while some additional domains are also present (Figures 1-7).




**Figure 1:** Conserved BRCA2 domains a) Homo sapiens b) Pan troglodytes c) Gorilla gorilla




**Figure 2:** Motif locations of BRCA2 gene with their p-values




**Figure 3:** Phylogenetic analysis of 25 species by using MEGA-7




**Figure 4:** Phylogenetic analysis of 25 species in circular form by using ClustalW 7



**Figure 5:** Phylogenetic analysis of candidate species by using MEGA-7



**Figure 6:** Positions of introns and exons are determined through gene structure display server, introns are marked by black colour while exons are marked by yellow colour.



**Figure 7:** Graphical map of BRCA2 by using serial cloner

**4.3. Gene Structure Analysis of BRCA2 Gene**

In evolutionary analysis, the locations of introns/exons in orthologous genes are particularly well-conserved, while in paralogous genes the location of introns and exons is found approximately less conserved. In order to confirm the structural variation of BRCA2 genes, we determined the positions of introns/exons by making comparison of CDS and genomic sequences. Although, there was substantial diversity found in the number and length of exons in all these candidate species including Homo sapiens, Pan troglodytes and Gorilla. Although, sequences retrieved at the subfamily level of these three species showed resemblance in gene structures in

respect to intron number, chromosome number and exon length. Conserved motifs of BRCA2 proteins were identified and proved useful to determine the protein functions, and the additional motifs of BRCA2 gene were found functional in activating the functions of BRCA2 proteins in humans (Lord & Ashworth, 2007) [13]. On 13 chromosome number BRCA2 gene is present in Homo sapiens. The distribution of BRCA2 gene was found dispersed on 13 chromosomes. The evolution of BRCA2 gene was further observed in all three mammalian species, while gene duplication events were examined for tandem and segmental duplications (Teul -Vega et al., 2015) [18] (Table-1).

**Table 1:** Proposed nomenclature and important features of BRCA2gene

Proposed Names	Gene Locus	Protein Accession #	RNA Accession #	Exon	Chr #	ORF length	Amino acid length	Start of Genomic Location	Conserved domains in protein sequence
BRCA2 in Homo sapiens	50	P51587.3	59.4	27	13	10257	3418	Chr13:32	BRCA-2_helical
BRCA2 Pan stroglydyt es	16780623	16780623.2	16925134.2	29	13	10257	3418	Chr13	BRCA-2_helical
BRCA2 in Gorilla	18895519	18895519.2	19039974.2	27	13	10167	3388	Chr13	BRCA-2_helical

**4.4. Mapping of BRCA2 Gene**

By using serial cloner, a graphical map of BRCA2 gene is obtained. By constructing a graphical map sequence length, nucleotides and base pairs are determined. Blunt ends are generated at 5 and 3 end. In serial cloner method protein sequence of BRCA2 in FASTA format is selected and mapping of gene has done. All genes are mapped with their respective restriction sites.

**5. Discussion**

This present study shows molecular analysis of BRCA2 gene in humans and animals. It has been observed that the patients who have mutations in BRCA2 gene are at greater risk of developing cancer than the noncarriers (Prakash, Zhang, Feng, & Jasin, 2015) [17]. Through current advancement in research scientists are able to edit genes by DNA repair mechanism via knock out and can directly determine the effects of mutation. Mutation carriers pres-

ent in genome provides valuable information about mechanism of mutation (Johnston et al., 2015) [10]. Different characteristics of mutant genes, sequence retrieved, conserved domains, positions of introns and exons are collectively meaningful to investigate relationship of *brca2* gene in mammalian species. Further, by using gene structure analysis exons count from 0 to 27 are examined precisely. In Pan troglodytes and Gorilla (western gorilla) closely related genes are observed because of their similarity in structure of introns and exons that shows their close phylogenetic relationship (Birkbak et al., 2013) [1-2]. Results of phylogenetic tree also depict that humans, chimpanzee and western gorilla share a common ancestor. The reason behind this hypothesis is similarity in their intron exon structure. On chromosome number 13, large number of BRCA2 genes are found in all three species. That shows there is same chromosomal map for human, chimpanzee and gorilla. Subsequently, random distribution of genes on same chromosome is a evidence of variation in genes. That results in a evolutionary process of three species which diverged from one another in their genome (Blokzijl, Janssen, Van Boxtel, & Cuppen, 2018) [3]. Through phylogenetic analysis, by comparing data in all three candidate species most closely related genes are observed. Hence, it is concluded that BRCA2 genes maintained their function even after divergence in humans, chimpanzee and western gorilla (Wiggins, Walker, & Pearson, 2020) [21].

## 6. Conclusion

This research showed a high similarity of BRCA2 in structure of intron and exons present in mammalian species. Although, duplication of genes in candidate species is clearly examined. This research study depict the close evolutionary relationship between humans and chimpanzee genes as well as with western gorilla. Through this study the identified relatable genes can be used as informational source to manipulate the mutations in BRCA2 gene that are fatal.

## 7. Conflicts of Interest

Author has no conflicts of interest.

## 8. Acknowledgment

Author is thankful to GCUF, for this study.

## References

1. Acedo A, Hernández-Moro C, Curiel-García Á, Díez-Gómez B, Velasco EA. Functional classification of BRCA2 DNA variants by splicing assays in a large minigene with 9 exons. *Human mutation*. 2015; 36(2): 210-221.
2. Birkbak NJ, Kochupurakkal B, Izarzugaza JM, Eklund AC, Li Y, Liu J, et al. Tumor mutation burden forecasts outcome in ovarian cancer with BRCA1 or BRCA2 mutations. *PloS one*. 2013; 8(11): e80023.
3. Blokzijl F, Janssen R, Van Boxtel R, Cuppen E. MutationalPatterns: comprehensive genome-wide analysis of mutational processes. *Genome medicine*. 2018; 10(1): 1-11.
4. Bolton KL, Chenevix-Trench G, Goh C, Sadetzki S, Ramus SJ, Karlan BY, et al. Association between BRCA1 and BRCA2 mutations and survival in women with invasive epithelial ovarian cancer. *Jama*. 2012; 307(4): 382-389.
5. Burke LJ, Sevcik J, Gambino G, Tudini E, Mucaki EJ, Shirley BC, et al. BRCA1 and BRCA2 5' noncoding region variants identified in breast cancer patients alter promoter activity and protein binding. *Human mutation*. 2018; 39(12): 2025-2039.
6. De Talhouet S, Peron J, Vuilleumier A, Friedlaender A, Viassolo V, Ayme A, et al. Clinical outcome of breast cancer in carriers of BRCA1 and BRCA2 mutations according to molecular subtypes. *Scientific reports*. 2020; 10(1): 1-9.
7. Eccles D, Mitchell G, Monteiro A, Schmutzler R, Couch F, Spurdle A, et al. BRCA1 and BRCA2 genetic testing—pitfalls and recommendations for managing variants of uncertain clinical significance. *Annals of oncology*. 2015; 26(10): 2057-2065.
8. Forbes C, Fayter D, De Kock S, Quek RG. A systematic review of international guidelines and recommendations for the genetic screening, diagnosis, genetic counseling, and treatment of BRCA-mutated breast cancer. *Cancer management and research*. 2019; 11: 2321.
9. Hatano Y, Tamada M, Matsuo M, Hara A. Molecular trajectory of BRCA1 and BRCA2 mutations. *Frontiers in Oncology*. 2020; 10.
10. Johnston JJ, Lewis KL, Ng D, Singh LN, Wynter J, Brewer C, et al. Individualized iterative phenotyping for genome-wide analysis of loss-of-function mutations. *The American Journal of Human Genetics*. 2015; 96(6): 913-925.
11. Jonsson P, Bandlamudi C, Cheng ML, Srinivasan P, Chavan SS, Friedman ND, et al. Tumour lineage shapes BRCA-mediated phenotypes. *Nature*. 2019; 571(7766): 576-579.
12. Kemp Z, Turnbull A, Yost S, Seal S, Mahamdallie S, Poyastro-Pearson E, et al. Evaluation of cancer-based criteria for use in mainstream BRCA1 and BRCA2 genetic testing in patients with breast cancer. *JAMA network open*. 2019; 2(5): e194428-e194428.
13. Lord CJ, Ashworth A. RAD51, BRCA2 and DNA repair: a partial resolution. *Nature structural & molecular biology*. 2017; 14(6): 461-462.
14. Lou DI, McBee RM, Le UQ, Stone AC, Wilkerson GK, Demogines AM, et al. Rapid evolution of BRCA1 and BRCA2 in humans and other primates. *BMC evolutionary biology*. 2014; 14(1): 1-13.
15. Peixoto A, Pinto P, Guerra J, Pinheiro M, Santos C, Pinto C, et al. Tumor Testing for Somatic and Germline BRCA1/BRCA2 Variants in Ovarian Cancer Patients in the Context of Strong Founder Effects. *Frontiers in Oncology*. 2020; 10: 1318.
16. Pfeiffer CM, Ho BN, Singh AT. The evolution, functions and applications of the breast cancer genes BRCA1 and BRCA2. *Cancer Genomics-Proteomics*. 2017; 14(5): 293-298.
17. Prakash R, Zhang Y, Feng W, Jasin M. Homologous recombination and human health: the roles of BRCA1, BRCA2, and associated proteins. *Cold Spring Harbor perspectives in biology*. 2015; 7(4): a016600.

18. Teulé-Vega À, Lázaro García C, Blanco Guillermo I, Valle Domínguez Jd. An original phylogenetic approach identified mitochondrial haplogroup T1a1 as inversely associated with breast cancer risk in BRCA2 mutation carriers. *Breast Cancer Research*. 2015; 17(61).
19. Théry JC, Krieger S, Gaildrat P, Révillion F, Buisine MP, Killian A, et al. Contribution of bioinformatics predictions and functional splicing assays to the interpretation of unclassified variants of the BRCA genes. *European Journal of Human Genetics*. 2011; 19(10): 1052- 1058.
20. Walker LC, Whiley PJ, Couch FJ, Farrugia DJ, Healey S, Eccles DM, et al. Detection of splicing aberrations caused by BRCA1 and BRCA2 sequence variants encoding missense substitutions: implications for prediction of pathogenicity. *Human mutation*. 2010; 31(6): E1484-E1505.
21. Wiggins GA, Walker LC, Pearson JF. Genome-wide gene expression analyses of BRCA1- and BRCA2-associated breast and ovarian tumours. *Cancers*. 2020; 12(10): 3015.
22. Zhao W, Wiese C, Kwon Y, Hromas R, Sung P. The BRCA tumor suppressor network in chromosome damage repair by homologous recombination. *Annual review of biochemistry*. 2019; 88: 221-245.