

## Primary Angiosarcoma of the Breast in an Adolescent Female

Cherie M Kuzmiak, DO, FACR, FSB<sup>†</sup>\* and Christopher Burns, DO\*

Department of Radiology, University of North Carolina, Chapel Hill, NC, USA

**\*Corresponding author:**

Cherie M Kuzmiak,  
DO, FACR, FSB, University of North Carolina,  
Department of Radiology, Chapel Hill, NC, USA

Received: 25 Mar 2026

Accepted: 06 Apr 2026

Published: 17 Apr 2026

J Short Name: Ajsccr

**Copyright:**

©2026 Cherie M Kuzmiak. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially

**Citation:**

Cherie M Kuzmiak, Primary Angiosarcoma of the Breast in an Adolescent Female. Ame journal of Sur and Clin Case Rep® 2026; V10(1): 1-4

### 1. Abstract

#### 1.1. History


A 17-year-old female presented to the emergency room with palpable right breast masses. On physical examination, there was diffuse ecchymosis of the right breast. A firm mobile mass was present in the upper outer and upper inner quadrant of the right breast. Due to the patient's age, a diagnostic US (Figure 1) was initially performed [1]. Afterwards, she underwent a diagnostic mammogram (Figure 2).

#### 1.2. Imaging Findings


Ultrasound of the right breast shows a 6.5 cm, irregular, heterogeneous mass with non-circumscribed margins and mixed posterior features within the upper outer quadrant. A 5.5 cm, oval, mixed solid and cystic mass, with circumscribed margins and parallel orientation is present within the upper inner quadrant (Figure 1). The mammogram demonstrates a 6.5 cm, irregular, isodense mass with indistinct margins within the upper outer

quadrant and a 5.5 cm, oval, isodense mass with indistinct margins in the upper inner quadrant (Figure 2). The left mammogram is normal (not shown). An US-guided biopsy of the two suspicious right breast masses were subsequently performed. Histopathology shows a primary angiosarcoma with high grade nuclear atypia and numerous mitotic figures at both sites (Figure 3). The patient then underwent a bilateral breast MRI with contrast.


The MRI demonstrates an asymmetrically enlarged right breast with two large masses that have heterogeneous T2 signal. The post-contrast images demonstrate a 6.5 cm, irregular mass with rim enhancement within the upper outer quadrant and a 5.5 cm, oval mass with rim enhancement in the upper inner quadrant of the right breast (Figure 4). The left breast is normal on MRI. The patient then underwent staging with F-18 FDG PET-CT, which shows increased heterogeneous rim uptake of the FDG involving the right breast masses (Figure 5). There was no distant disease.




**Figure 1:** Large field-of-view ultrasound of a 17-year-old female with palpable right breast masses. The image shows a 6.5 cm, irregular, heterogeneous mass (arrowheads) with non-circumscribed margins and mixed posterior features with shadowing (^) within the upper outer quadrant. In addition, a 5.5 cm, oval, mixed solid and cystic (\*) mass (arrows), with circumscribed margins and parallel orientation is present within the upper inner quadrant.




**Figure 2:** Mammographic images of a 17-year-old female with palpable right breast masses. Mediolateral oblique (A) and craniocaudal (B) mammogram images demonstrate a 6.5 cm, irregular, isodense mass with indistinct margins within the upper outer quadrant of the breast (arrowhead). A 5.5 cm, oval, isodense mass with indistinct margins is also present in the upper inner quadrant of the breast (arrow). The left mammogram is normal (not shown).



**Figure 3:** Histopathology of the right breast. Primary angiosarcoma with high-grade nuclear atypia (arrows) is present. Hematoxylin and eosin stain, 40x.



**Figure 4:** Bilateral breast MRI of a 17-year-old female with palpable right breast masses. Pre-contrast non-fat suppression axial T2 (A) and axial T1 fat suppression (B) demonstrate an asymmetrically enlarged right breast with two large masses (arrows) that have heterogeneous T2. The post-contrast axial T1 image (C) and post-contrast axial T1 subtraction image (D) demonstrate a 6.5 cm, irregular mass with rim enhancement (arrowhead) within the upper outer quadrant and a 5.5 cm, oval mass with rim enhancement (arrow) in the upper inner quadrant of the right breast. No nodal involvement is seen. The left breast is normal.



**Figure 5:** Axial fused F-18 FDG PET-CT demonstrates increased heterogeneous rim uptake of FDG involving the right breast masses (arrows). There was no evidence of metastatic disease.

### 1.3. Differential Diagnosis

The differential diagnosis for a mixed solid and cystic mass in a young female patient include infection and hematoma/trauma in the appropriate clinical setting. Less common etiologies to consider include an infarcted fibroadenoma, papilloma, phyllodes tumour, invasive ductal cancer and primary breast angiosarcoma.

### 1.4. Discussion

Primary angiosarcoma (PAS) is a malignancy of breast endothelial origin that is not related to prior radiation and has an incidence of 0.04% [2,3]. It typically presents in younger women as a palpable mass and 20% may have overlying skin changes,

bluish discoloration [2,4-6].

The imaging findings of PAS are nonspecific. On mammography, it may present as a noncalcified mass with circumscribed or indistinct margins [2,4-6]. Thirty-three percent of PASs have been reported to be occult except for minor skin thickening [2,6]. The sonographic ranges from a hypoechoic to heterogeneous mass with circumscribed or not circumscribed margins to a region of mixed echogenicity [2,4-6]. PAS may be hyper vascular on Power Doppler [2,6]. On MR, it may have high-signal intensity on T2-weighted images with variable kinetics depending on tumour grade [2,4-6]. Due to its nonspecific appearance, needle core biopsy is needed for a definitive diagnosis.

The treatment of PAS is surgical resection with negative margins [2,6]. Nodal involvement is rare since metastases are hematogenous [2,6]. Neoadjuvant or adjuvant chemoradiation therapy have been proposed [2, 6]. The reported overall 5-year survival rate for PAS is 49% [7].

The patient underwent neoadjuvant chemoradiation therapy followed by a right mastectomy. Final histopathology showed a 12.2 cm PAS.

### References

1. Katherine A Klein, Maddi Kocher, Andrea Porpiglia, MD, Expert Panel on Breast Imaging; Klein K, Kocher M. ACR Appropriateness Criteria® Palpable Breast Masses: 2022 Update. *J Am Coll Radiol.* 2023; 20(5S): S146-S163.
2. Glazebrook K, Magut M, Reynolds C. Angiosarcoma of the breast. *AJR Am J Roentgenol.* 2008; 190(2): 533-38.
3. Young RJ, Brown NJ, Reed MW. Angiosarcoma. *The Lancet Oncology.* 2010; 11(10): 983-991.
4. Yang WT, Hennessy BT, Dryden MJ, Valero V. Mammary angiosarcomas: imaging findings in 24 patients. *Radiology.* 2007; 242: 725-734.
5. Kapoor M, Yoon E, Yang W, Patel M. Breast Angiosarcoma: Imaging Features with Histopathologic Correlation, *Journal of Breast Imaging.* 2023; 5(3): 329-338.
6. Friedrich A, Reisenbichler E, Heller D. Characteristics and long-term risk of breast angiosarcoma. *Ann Surg Oncol.* 2021; 28(9): 5112-5118.